



Figures Reblissing for:
***Search for Heavy, Long-Lived
Neutralinos that Decay to Photons at
CDFII using Photon Timing***
(to be submitted to PRD)

**Paul Geffert, Max Goncharov, Slava
Krutelyov*, Eunsin Lee, Rishi Patel**,
David Toback and Peter Wagner*****

*Now postdoc at UCSB

** REU student from NYU

*** Now postdoc at Penn



Why we're re blessing

- Delayed Photon Analysis published in PRL
 - PRL 99, 121801 (2007)
- Publish more details in a full PRD
 - Paper Seminar today
- Godparents: F. Bedeschi, H. Budd, A. Messina
- Today: Current versions of figures updated since CDF notes were posted
 - CDF Note 9171
- *No New Results*



Outline

- Short Overview of the Motivation and Theory
- Brief Summary of the Analysis
- Plots
- Conclusion

Supporting Documentation: CDF Notes 7515, 7918, 7928, 7929, 7960, 8015, 8016



Motivation and Theory


- GMSB models predict heavy neutralinos that decay to photons (see next slide)
- $ee\gamma\gamma + \cancel{E}_T$ candidate event at CDF in Run I
- First search for heavy, long-lived particles that decay to photons at a hadron collider



GMSB Models

- The lightest neutralino is the NLSP and decays into a gravitino and a photon
- For much of the parameter space the neutralino decay time can be \sim ns
- At the Tevatron neutralinos are pair produced from $\tilde{\chi}_1^+ \tilde{\chi}_1^-$ or $\tilde{\chi}_1^+ \tilde{\chi}_2^0$

Brief Summary of the Analysis



Signature

$\Rightarrow \cancel{E}_T$

Jets

$\Rightarrow \gamma\gamma$ or γ

- $\gamma + \cancel{E}_T + \text{jets}$ analysis is sensitive to ns lifetimes while $\gamma\gamma + \cancel{E}_T$ analysis is sensitive to prompt neutralino decays


Toback and Wagner, PRD 70, 114032 (2004)



Next



*Tell the story of the analysis
using the PRD figures...*

PRD Figure 1- Feynman Diagrams




- Feynman diagrams of the dominant production processes at the Tevatron
- Use SPS 8 GMSB model line (Eur. Phys. J. C 25, 113 (2002)): $\tan(\beta)=15$, $\text{sgn}(\mu)=1$, $N_m=1$, and $M_m=2\Lambda$

PRD Figure 2- Event Schematic and Time Distribution



- Left- Schematics of a long-lived neutralino decay into a gravitino and a photon
- Right- The corrected time distribution for a GMSB example point as well as the non-collision and SM backgrounds

Fig. 3- Photon Can Hit the Calorimeter with a Large Incident Angle



- Left- Definition of α , the projection of the photon incident angle in the (r, z) plane
- Right- Definition of β , the projection of the photon incident angle in the (r, ϕ) plane

Fig. 4- Look at Photon Incident Angles

- Delayed photons have a larger incident angle than promptly produced photons
- Distribution of the total incident angle, ψ , of the photon at the face of the calorimeter







Fig. 5- Compare ID Variables (Long Lifetime vs. 0 Lifetime)



CDFSim ID variable distributions minus their requirement value

Fig. 6- Efficiency vs. Angle, Compare Electrons and Photons from Data and MC



- Left- The efficiencies for e's and γ 's to pass ID requirements vs. incident angle α
- Right- The same but for β
 - Efficiency falls in β primarily due to the energy isolation requirement; small effect, well-modeled in MC

Fig. 7- New PMT Asymmetry Cut to Kill Spikes

- Compare asymmetry of spikes to real electrons
- Require asymmetry to be less than 0.6






Fig. 8- Vertexing



- The collision time and position for the reconstructed highest Σp_T vertex in $W \rightarrow e\nu$ events
- Also show correlation for fun







Fig. 9- Vertexing Performance/ Resolution




The difference in z and t between two arbitrarily selected sets of tracks from the same reconstructed vertex in a $W \rightarrow e\nu$ dataset

Fig. 10- Vertexing Performance continued; Compare Vertex to Electron Track



- The distributions are centered at zero → no clustering bias
- The second Gaussian contains events where the electron is from a second vertex in the event

Fig. 11- Vertexing Efficiency



We require a vertex to have at least 4 tracks and $\Sigma p_T > 15$ GeV \rightarrow 100% efficiency

Fig. 12- Check EMTiming Simulation and Show Resolution

Well centered around 0 ns with RMS of 0.64 ns








Fig. 13- Timing Distribution for “Right” and “Wrong” Vertex Selection





- Top Left- Electron track matches vertex (“Right Vertex”)
- Top Right- Electron anti-matched to vertex (“Wrong Vertex”)

Fig. 14- Systematic Variation of Timing Mean and RMS



Look at the timing distribution for electrons from subsamples of $W \rightarrow e\nu + \text{jets}$ events for different requirements on electron E_T , jet E_T , and E_T

Fig. 15- Systematic Variation of Timing – Wrong Vertex



The mean and RMS of the timing for electrons as a function of η , when the wrong vertex is picked

Fig. 16- Beam Halo

- **Illustration of a beam halo event**
- **Top-**The mean corrected time changes as a function of η but is always less than zero
- **Bottom-**The halo interacts with many hadronic calorimeter towers at high η




Fig. 17- Beam Halo vs. Cosmics

The variables used to separate cosmic and beam halo backgrounds to create their timing templates





Fig. 18- Timing for Beam Halo and Cosmics



The corrected time distributions for beam halo (left) and cosmic ray (right) backgrounds

Fig. 19- More on Beam Halo

- **Most beam halo photons arrive at $\phi \approx 0$**
- **Use this for background normalization**




Fig. 20- Analysis Optimization

The expected 95% C.L. cross section limit as a function of the lower value of the timing requirement for a GMSB example point





Fig. 21- The Data... Timing Distribution



- Left- The timing distribution for the signal and all backgrounds
- Right- A zoomed in view of the signal region, [2,10] ns
- Two events are observed in the signal region, consistent with the background expectation of 1.3 ± 0.7 events

Fig. 22- Kinematic Distributions

- Compare background predictions and data
- No evidence for new physics









Fig. 23- Expected and Observed Limits and Production Cross Sections



- Limits vs. lifetime for $m=100$ GeV
- Limits vs. mass for a lifetime of 5 ns

Fig. 24- Results...

The contours of constant 95% C.L. cross section upper limit for the observed number of events




Fig. 25- Exclusion Region

- Expected and observed 95% C.L. exclusion region along with LEP limits
- Highest mass reach is 108 GeV (expected) and 101 GeV (observed) for a lifetime of 5 ns.





Fig. 26- Expected Sensitivity for 2 fb⁻¹ and 10 fb⁻¹

- Background scaled with luminosity
- The shaded band shows the cosmologically favored parameter space





Conclusion

- Both readings of the PRD are complete and the GPS have signed off
- Paper seminar today at the Weekly meeting